

Production of 2,3-Dihydroxybenzoic Acid by *Brucella* Species

I. López-Goñi, I. Moriyón

Departamento de Microbiología. Facultad de Medicina. Universidad de Navarra.

Abstract of:

Current Microbiology Vol. 31 (1995), pp. 291-293

A total of 29 strains of *Brucella abortus*, *B. melitensis*, *B. suis*, *B. ovis*, and *B. neotomae* were examined for growth and catechol production in a semisynthetic low-iron medium. All strains showed reduced growth yields and, quantitatively, production of catechols va-

ried widely among the different strains with no relationship to species, biotypes, or serotypes or *Bruce-lla*. No clear correlation between catechol productions and growth under iron-limiting conditions was observed. The major catechol was identified as 2,3-dihydroxybenzoic acid, and neither other iron-regulated catechols nor hydroxamate type compounds were detected when representative strains of *B. Abortus melitensis* were grown in tryptic soy broth in the presence of iron-sequestering agents.

Evaluation of PCR and Indirect Enzyme-Linked Immunosorbent Assay on Milk Samples for Diagnosis of Brucellosis in Dairy Cattle

C. Romero, M. Pardo, M.J. Grillo, R. Díaz, J.M. Blasco, and I. López-Goñi

Departamento de Microbiología, Universidad de Navarra, and
Unidad de Sanidad Animal, Servicio de Investigación Agraria, Diputación General de Aragón, Zaragoza, Spain.

Abstract of:

Journal of Clinical Microbiology, Dec. 1995,
p. 3198-3200

A study was performed to evaluate the previously described PCR (C. Romero, C. Gamazo, M. Pardo, and I. López-Goñi, J. Clin. Microbiol. 33:615-617, 1995) for the diagnosis of brucellosis in dairy cattle. Milk samples from 56 *Brucella* milk culture-positive cattle and from 37 cattle from *Brucella*-free herds were examined for *Brucella* DNA by PCR and for specific antibodies by an indirect enzyme-linked immunosorbent as-

say (ELISA). The specificities of both tests were 100 % when testing the milk samples from *Brucella*-free cattle. The milk samples from 49 infected cattle were positive by PCR (87.5 % sensitivity), and 55 were positive by ELISA (98.2 % sensitivity). A PCR-positive sample was negative by ELISA, and 7 ELISA-positive samples were PCR negative, yielding an observed proportion of agreement of 0.91 for the two tests. Although the results suggest that ELISA is a better screening test than PCR, the combined sensitivity of the two assays was 100 %, and their simultaneous applications could be more useful than one test alone for a rapid screening of brucellosis in dairy cattle.